

# MIDTERM REVIEW


## Observational Evidence for mass discrepancies

- Oort discrepancy in solar neighborhood
- flat rotation curves of spiral galaxies
- clusters of galaxies
  - velocity dispersions
  - hydrostatic equilibrium of X-ray gas
  - gravitational lensing of background galaxies
- large scale structure
- $\Omega_m > \Omega_b$

## Early indications included

- Oort (1932) - factor of  $\sim 2$  discrepancy in solar neighborhood
- Zwicky (1933) - factor of  $\sim 100$  discrepancy in clusters
- Ostriker & Peebles (1973) - factor of  $\sim 10$  discrepancy in bar instability in disks

## Dark Matter Candidates



WIMPs favored because

- $\Omega_m > \Omega_b$
- LSS formation

Dark matter is the usual inference; the discrepancy might also indicate a change in dynamical laws

## Virial Theorem

Can be derived from stationary moment of inertial tensor

$$\text{boils down to } 2\langle K \rangle + \langle W \rangle = 0$$

Kinetic E      Potential Energy

for  $N$  particles of equal mass  $m$  such that  $M = Nm$ ,

$$M = \frac{2\sigma^2 R_{\text{rms}}}{G} \quad \text{where the harmonic radius } R_{\text{rms}}$$

is usually approximated as  $R_{\text{rms}} \approx 1.25 R_d$

Vertical Force (Oort: restoring force to disk)

$$K_z = -\frac{\partial \Phi}{\partial z} = \frac{1}{v} \frac{\partial (v\sigma^2)}{\partial z} \quad \text{where } v(z) \text{ is the vertical profile of tracer population}$$

locally this boils down to

$$\sigma_z^2 = 2\pi G \Sigma z_0$$

e.g.  $v(z) = v_0 e^{-z/z_0}$

## Disk Stability

LOCAL: Toomre  $Q$ :  $Q = \frac{v R K}{3.36 G \Sigma}$  locally stable if  $Q \geq 1$

GLOBAL:  $\chi_m = \frac{K^2 R}{2\pi m G \Sigma}$  higher surface densities less stable

Ostriker & Peebles:  $t \lesssim 0.14$  where  $t = \frac{T}{|W|}$

with

$$K = T + \frac{1}{2}\Pi \quad T = \text{rotational kinetic energy}$$
$$\Pi = \text{kinetic energy in random motions}$$

## Exponential Disks

2D face on surface brightness profile

$$\Sigma(R) = \Sigma_0 e^{-R/R_d}$$

$\Sigma_0$  = central surface density  
 $R_d$  = scale length

This integrates to a total luminosity  $L = 2\pi \Sigma_0 R_d^2$

The enclosed luminosity is a simple fun of ~~the~~ scale length

$$L(<x) = 2\pi \Sigma_0 R_d^2 \left[ 1 - (1+x)e^{-x} \right] \quad \text{where } x = \frac{R}{R_d}$$

in 3D one can have a "double exponential" model

$$\rho = \rho_0 e^{-R/R_d} e^{-z/z_0}$$

For general light profiles we can fit the

Sersic profile

$$\Sigma(R) = \Sigma_e e^{-b_n \left[ \left( \frac{R}{R_e} \right)^{1/n} - 1 \right]}$$

which reduces to the exponential form for  $n=1$   
and is equivalent to the de Vaucouleurs profile for  $n=4$

## Potential - Density Pairs

Poisson Equ  $\nabla^2 \Phi = 4\pi G \rho$

has an analytic solution for a handful of  $\Phi$ - $\rho$  pairs.

$\therefore$  It helps to know  $\nabla^2$  in the right coordinate system

Phase space  $f(\vec{x}, \vec{v}, t)$  : the distribution function

describes the density distribution of particles in both configuration space  $(x, y, z)$  and momentum  $(v_x, v_y, v_z)$ .

For a stationary (stable) system of widely separated stars, we have the collisionless Boltzmann eqn

$$\frac{\partial f}{\partial t} + \vec{v} \cdot \vec{\nabla} f - \vec{\nabla} \Phi \cdot \frac{\partial f}{\partial \vec{v}} = 0$$

Note that the density in configuration space  $\nu$  is just the integral over  $f$  in momentum:

$$\nu = \int f d^3v$$

We usually use this to break the collisionless Boltzmann eqn into the

Jean's eqns

$$\frac{\partial \nu}{\partial t} + \frac{\partial (\nu \bar{v}_i)}{\partial x_i} = 0$$

$$\frac{\partial (\nu \bar{v}_j)}{\partial t} + \frac{\partial (\nu \overline{v_i v_j})}{\partial x_i} + \nu \frac{\partial \Phi}{\partial x_j} = 0$$

$$\nu \frac{\partial \bar{v}_j}{\partial t} + \nu \bar{v}_i \frac{\partial \bar{v}_j}{\partial x_i} = -\nu \frac{\partial \Phi}{\partial x_j} - \frac{\partial (\nu \sigma_{ij}^2)}{\partial x_i}$$

where we have the "moments" of  $f$

$$\nu = \int f d^3v$$

$$\overline{v_i v_j} = \frac{1}{\nu} \int v_i v_j f d^3v$$

$$\bar{v}_i = \frac{1}{\nu} \int v_i f d^3v$$

$$\sigma_{ij}^2 = \overline{v_i v_j} - \bar{v}_i \bar{v}_j$$

## Energy & Angular momentum



$$E = \frac{1}{2} (v_R^2 + v_\phi^2 + v_z^2) + \Phi(R, z) \quad \text{energy per unit mass}$$

in cylindrical coordinates

$$J_z = R v_\phi \quad \text{angular momentum per unit mass}$$

## Effective Potential

$$\Psi(R, z) = \Phi(R, z) + \frac{J_z^2}{2R^2}$$



## Galactic constants

$R_0$  - distance to Galactic Center

$V_0$  - circular velocity of LSR (sometimes called  $\Theta_0$ )

$\Omega_0 = \frac{V_0}{R_0}$  - orbital frequency

0 subscript denotes solar location

orbital period

$$P = \frac{2\pi}{\Omega}$$

## Oort "constants"

$$A = -\frac{1}{2} \left[ R \frac{d\Omega}{dR} \right]_{R_0} = \frac{1}{2} \left( \frac{V}{R} - \frac{dV}{dR} \right)_{R_0} \quad \text{SHEAR}$$

$$B = \frac{1}{2} \left( \frac{V}{R} + \frac{dV}{dR} \right)_{R_0} \quad \text{VORTICITY}$$

due to

angular

momentum

gradient

NOTE :  $\Omega = A - B$

$$-\frac{dV}{dR} = A + B$$

epicyclic frequency :  $K^2 = -4B\Omega$


frequencies  $\omega_z > K > \Omega$  so orbits not closed

more vertical oscillations than radial than complete orbits

$$\omega_z \approx 48$$

$$K \approx 37$$

$$\Omega \approx 30 \text{ km s}^{-1} \text{ kpc}^{-1} \text{ in solar neighborhood}$$



size of ellipsoid

$$\frac{Y}{X} = \frac{2\Omega}{K} = \frac{\sigma_y}{\sigma_x}$$

timescales

crossing time  $t_c = \frac{2R}{V}$

dynamical time  $t_d = \sqrt{\frac{3\pi}{16G\rho}}$

relaxation time  $\frac{t_r}{t_c} = \frac{N}{48f^2} \approx \frac{N}{6 \ln(N/2)}$

### 3 LAWS of GALACTIC ROTATION

#### 1. Flat rotation curves

The rotation curves of rotating galaxies tend to approach an approximately constant velocity that persists to indefinitely large radii.

#### 2. Baryonic Tully-Fisher Relation: $M_b = AV_f^4$

The total baryonic mass of a rotating galaxy scales as the fourth power of its flat rotation velocity.

#### 3. Mass Discrepancy - Surface Density relation

The amplitude of the mass discrepancy scales roughly as  $\Sigma_b^{1/2}$ .  
(Holds both globally and locally)

## Halo models

pseudo-isothermal

empirically motivated

characterized by

core radius  $R_c$

flat velocity  $V_{\infty}$

$$\rho_{\text{iso}}(r) = \frac{\rho_0}{1 + (r/R_c)^2}$$

$$V(r) = V_{\infty} \sqrt{1 - \frac{R_c}{R} \tan^{-1}\left(\frac{R}{R_c}\right)}$$

$$V_{\infty} = \sqrt{4\pi G \rho_0 R_c^2}$$

## NFW

derived from simulations

characterized by

$c$

$R_{200}/V_{200}/M_{200}$

$$\rho_{\text{NFW}}(r) = \frac{4\rho_s}{(r/r_s)(1+r/r_s)^2}$$

$$V(r) = V_{200} \sqrt{\frac{\ln(1+cx) - \frac{cx}{1+cx}}{x \left[ \ln(1+c) - \frac{c}{1+c} \right]}}$$

$$x = \frac{r}{R_{200}}$$

$$\frac{r}{R_{200}}$$

$$c = \frac{R_{200}}{r_s}$$

$$V_{200} = h R_{200}$$

in km/s                  in kpc

where  $h = \frac{H_0}{100}$

$$M_{200} = \frac{4\pi}{3} (200) R_{200}^3$$

see also Einasto  
Burkert




## Cosmology essentials

Friedmann eqn:  $\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho - \frac{kc^2}{a^2} + \frac{\Lambda}{3}$

$$H = \frac{\dot{a}}{a} \quad \Omega_m = \frac{\rho}{\rho_{\text{crit}}} \quad \rho_{\text{crit}} = \frac{3H^2}{8\pi G}$$

In the early universe, regions with local  $\Omega_m > 1$   $\left(\frac{\delta\rho}{\rho} > 1\right)$   
can begin to gravitationally collapse  
IFF they are composed of a form of  
dark matter that does not interact with photons

Monolithic galaxy formation  
replaced by hierarchical galaxy formation



Overdensities

$$M_\Delta = \frac{4\pi}{3} \Delta \rho_{\text{crit}} R_\Delta^3$$

$$V_\Delta = \frac{GM_\Delta}{R_\Delta}$$

$$\therefore M_\Delta = \frac{\Delta}{2G} H_0^2 R_\Delta^3 = \sqrt{\frac{2}{\Delta}} \frac{V_\Delta^3}{GH_0}$$