


Extrasolar Planets

A futuristic landscape with a large reddish planet in the sky and two bright stars. The foreground is a rocky, orange-brown terrain with jagged mountains in the background. The sky is dark blue with many small stars.


- **Today**
 - **Dwarf Planets**
 - **Extrasolar Planets**
- **Next week**
 - **Review Tuesday**
 - **Exam Thursday**
 - **also, Homework 6 Due**
 - **will count best 5 of 6 homeworks**

Dwarf Planets

- Large Kuiper Belt objects like Pluto now considered “dwarf planets.”
- Many now known




Dwarf planets of the Kuiper belt




There are many, many smaller (but still large) bodies in the Kuiper belt

Dwarf Planets - there's a lot out there!




Many of the recently discovered dwarf planets have orbits on one side of the solar system. This leads to a hypothesized massive Planet 9 that their orbits are in resonance with.




- Direct: pictures or spectra of the planets themselves
- Indirect: measurements of stellar properties revealing the effects of orbiting planets
 - Astrometric method (face-on)
 - Doppler method (edge-on)
 - Transit method (edge-on)
 - Gravitational lensing

Direct Imaging





Coronagraph blocks light from star, which is otherwise overwhelming. Even blocked, a tiny fraction of scattered light make the mess you see here.

Astrometric Technique



- We can detect planets by measuring the change in a star's position on sky.
- However, these tiny motions are very difficult to measure (~ 0.001 arcsecond).
- Best seen face-on


Doppler Technique



a A periodic Doppler shift in the spectrum of the star 51 Pegasi shows the presence of a large planet with an orbital period of about 4 days. Dots are actual data points; bars through dots represent measurement uncertainty.

- Best seen edge-on

Transit Technique




- A transit is when a planet crosses in front of a star.
- The resulting eclipse reduces the star's apparent brightness and tells us planet's radius.
 - best seen edge-on

Gravitational lensing Technique


Gravitation Microlensing




Extrasolar planet detected by gravitational microlensing




Exoplanet discoveries up to 2014



All methods
have strong
selection
effects






...and are good at measuring different things

Kepler is excellent at measuring radius; not necessarily mass.

What properties of extrasolar planets can we measure?




Measurable Properties


- Orbital period, distance, and shape (eccentricity)
- Planet mass, size, and density
- Atmospheric properties (sometimes)

*Exoplanet app
gives you scientific data
for thousands of exoplanets*

What can Doppler shifts tell us?




For a more massive planet in a similar orbit, we observe a larger Doppler shift with the same period.



- Doppler shift data tell us about a planet's mass and the shape of its orbit.

Planet Mass and Orbit Tilt




b We can detect a Doppler shift only if some part of the orbital velocity is directed toward or away from us. The more an orbit is tilted toward edge-on, the greater the shift we observe.

- We cannot measure an exact mass for a planet without knowing the tilt of its orbit, because Doppler shift tells us only the velocity toward or away from us.
- Doppler data give us lower limits on masses, $M \cdot \sin(i)$

Relative
Radii

From transits



*The Relative Sizes of
Known Exoplanets and **Solar System Planets***


Alex H. Parker / @Alex_Parker
Data from <http://exoplanet.eu/>

Astronomy Picture of the Day

[over the cosmos!](#) Each day a different image or photograph of our fascinating universe is featured, along with a brief explanation written by a professional astronomer.



<https://apod.nasa.gov/apod/ap151205.html>

2015 December 5




Kepler Orrery IV

The Kepler 11 system



- The periods and sizes of Kepler 11's 6 known planets can be determined using transit data.
- These periods are short
 - longest Kepler g at 118 days
- Tightly packed system!


Kepler-11 System



Venus


Mercury

Solar System



Calculating density


- Using mass, determined using the Doppler technique, and size, determined using the transit technique, density can be calculated.



For transiting planets, the Doppler method gives an accurate mass.

planet density:


$$\frac{\text{mass}}{\text{volume}} = 8.8 \text{ g/cm}^3$$



The transit method yields a radius, from which we can calculate the planet's volume.


The Kepler 11 system

- The densities of all these planets are low
 - 0.6 - 1.7 g/cc (Typical of Jovian planets)
- What does this imply about the solar nebula hypothesis?




The Kepler 11 system

- The densities of all these planets are low
 - 0.6 - 1.7 g/cc
- Star is
 - 0.96 mass of sun
 - 1.07 radius of sun
 - 8.5 Gyr old (sun is 4.5 Gyr)
- Tightly packed system that is nevertheless stable
 - despite lack of orbital resonances




Planetary Properties




Exoplanet properties in general: size vs. mass

Radius



Might exoplanets be home to life?




Habitable Zone

for terrestrial life

- Depends on
 - Brightness of star
 - also the spectrum of star - too much UV? too little?
 - Distance of planet from star
 - Nature of planet
 - surface gravity
 - atmosphere
 - greenhouse gases
 - water
 - other... some argue the moon is a necessary shield against too many major impacts

Presumably, the habitable zone is farther out from hot, bright stars and closer in to faint, cool ones.





Stars evolve over billions of years, gradually becoming brighter - planets may slip out of the habitable zone as a result

For the Earth, expect

- The sun increases slowly in brightness
- in 600 million years, the Carbon cycle may have progressed to the point that the atmosphere may lack sufficient CO₂ to sustain plant life
- In ~ 1 billion years, solar luminosity will have increased ~10%, evaporating the oceans (“wet greenhouse”)
- In ~6 billion years, the sun will expand into a red giant, potentially swelling far enough to encompass the orbit of the Earth

Future Earth?





GAME OVER MAN! GAME OVER!!

quickmeme.com