

Article

Tidal Dwarf Galaxies: Disc Formation at $z \simeq 0$

Federico Lelli ^{1,*}, Pierre-Alain Duc ², Elias Brinks ³ and Stacy S. McGaugh ¹

¹ Department of Astronomy, Case Western Reserve University, 10900 Euclid Ave, Cleveland, OH 44106, E-Mail: stacy.mcgough@case.edu

² Laboratoire AIM, CNRS, CEA/DSM, Université Paris Diderot, F-91191 Gif-sur-Yvette, France; E-Mail: paduc@cea.fr

³ Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK; E-Mail: e.brinks@herts.ac.uk

* Author to whom correspondence should be addressed; E-Mail: federico.elli@case.edu.

Academic Editor: Alexei Moiseev, José Alfonso López Aguerra and Enrichetta Iodice

Received: 15 October 2015 / Accepted: 2 November 2015 / Published: 5 November 2015

Abstract: Collisional debris around interacting and post-interacting galaxies often display condensations of gas and young stars that can potentially form gravitationally bound objects: Tidal Dwarf Galaxies (TDGs). We summarise recent results on TDGs, which are originally published in Lelli *et al.* (2015, A&A). We study a sample of six TDGs around three different interacting systems, using high-resolution HI observations from the Very Large Array. We find that the HI emission associated to TDGs can be described by rotating disc models. These discs, however, would have undergone less than one orbit since the time of the TDG formation, raising the question of whether they are in dynamical equilibrium. Assuming that TDGs are in dynamical equilibrium, we find that the ratio of dynamical mass to baryonic mass is consistent with one, implying that TDGs are devoid of dark matter. This is in line with the results of numerical simulations where tidal forces effectively segregate dark matter in the halo from baryonic matter in the disc, which ends up forming tidal tails and TDGs.

Keywords: interacting galaxies; peculiar galaxies

1. Introduction

Collisional debris around interacting and post-interacting galaxies are often observed to contain stellar and gaseous condensations that have masses, sizes, and star-formation rates (SFRs) comparable to those

of dwarf galaxies [1,2]. Zwicky [3] was the first to suggest that these condensations may collapse under self-gravity and form new independent entities: tidal dwarf galaxies (TDGs). Numerical simulations of interacting galaxies predict the formation of TDGs [4,5] and suggest that they may survive for several Gyr against internal or external disruption [6]. Interestingly, these simulations predict that TDGs should be free of non-baryonic dark matter (DM). This occurs because of two basic dynamical effects [7]: (i) tidal forces effectively segregate baryons in the disc (which is dynamically cold and can form tails and TDGs) from DM in the halo (which is too dynamically hot to form narrow tails); and (ii) once a TDG has formed, it has a shallow potential well and cannot accrete DM particles with typical velocity dispersions of $\sim 200 \text{ km s}^{-1}$. Similarly, TDGs are moving at the characteristic speed of the central DM halo ($\sim 200 \text{ km s}^{-1}$) and cannot be accreted by DM sub-haloes that have shallow potential wells ($\sim 50 \text{ km s}^{-1}$).

Bournaud *et al.* [8] used high-resolution HI observations to study the dynamics of three TDGs around NGC 5291. Puzzlingly, they found that the dynamical masses (M_{dyn}) of these TDGs are a factor of ~ 3 higher than the baryonic masses (M_{bar}). They hypothesized that the missing mass may consist of very cold molecular gas, which is not traced by CO observations but may constitute a large portion of baryonic DM in galaxy discs [9]. Alternatively, it was shown that MODified Newtonian Dynamics (MOND) can reproduce the observed rotation velocities of these TDGs without any need of DM [10,11]. In this regard, TDGs may help to distinguish between MOND and the standard model of cosmology, given that these two paradigms predict different internal dynamics for these recycled galaxies [12].




Figure 1. Interacting systems with genuine Tidal Dwarf Galaxies (TDGs, indicated in yellow). Optical images are overlaid with HI emission from the VLA (blue) and FUV emission from GALEX (pink), tracing star-forming regions. The red bar corresponds to ~ 10 kpc for the assumed distances.

Here I summarize recent results on the internal dynamics of TDGs (published in Lelli *et al.* [13]), which are based on high-resolution HI observations from the Very Large Array. We investigate six TDGs around three different interacting systems (Figure 1): the late-stage merger NGC 7252 (“Atoms

for Peace”), the collisional galaxy NGC 5291, and the post-merger lenticular NGC 4694. In particular, we revisit the DM content estimated by Bournaud *et al.* [8] for NGC 5291. We consider only *genuine* TDGs that satisfy the following requirements: (i) they are real condensations of gas and stars, which are not due to projection effects along the debris; (ii) they have higher metallicities than galaxies of similar mass, indicating that they are not pre-existing dwarf galaxies but recycled objects forming out of pre-enriched gas [13,14]; and (iii) they display a velocity gradient that is kinematically decoupled from the tidal debris, pointing to a local potential well [8,13].

2. Dynamics of TDGs

In Lelli *et al.* [13], we study the HI kinematics of TDGs by building 3D kinematical models. These techniques have been widely used and validated for typical dwarf galaxies [15,16]. In short, we construct disc models (with given radius, rotation velocity, and velocity dispersion) and project them on the sky to obtain model cubes that can be directly compared with the observed cubes. For all TDGs, we find that rotating disc models can reproduce the observed HI kinematics. An example (NGC 7252NW) is presented in Figure 2. The top panels show a *R*-band image (left), the HI map (middle), and the HI velocity field (right). Clearly, the gas distribution and kinematics resemble a rotating disc. The bottom panels show position-velocity diagrams obtained from the observed (left), model (middle), and residual (right) cubes along the major and minor axes. A disc model with axisymmetric kinematics can reproduce both the major and minor axes, suggesting that possible non-circular motions are relatively small. These techniques represent a major improvement over previous kinematic studies of TDGs because they allow us to estimate rotation velocities considering the effects of the gas distribution, velocity dispersion, disc inclination, and instrumental resolution. In particular, we find that the rotation velocities from [8] were over-estimated because the line-broadening due to turbulent motions was not taken into account.

Although a rotating disc provides a good description of the HI kinematics, there is an important issue with this model. The orbital times at the edge of the disc are longer than the dynamical timescale for the interaction/merger event estimated from numerical simulations. Hence, these HI discs have undergone less than a full rotation since the epoch of the TDG formation, raising the question of whether they are in dynamical equilibrium. For collisionless stellar systems, the process of relaxation takes a few orbital times due to the energy exchange between individual particles and the total gravitational potential [17]. For dissipative gas systems as TDGs, the energy can probably be exchanged and radiated away on a local crossing time (~ 10 Myr), thus one may speculate that the process of relaxation in collisional systems is faster than in collisionless ones. This remains an open issue that deserves further investigation.

If we assume that TDGs are in equilibrium, we can estimate dynamical masses using the observed rotation velocities (after correction for pressure support). We also estimate baryonic masses as the sum of atomic gas ($1.33 M_{\text{HI}}$), molecular gas (M_{mol}), and stars (M_*). In TDGs M_{bar} is heavily dominated by atomic gas, hence typical uncertainties on M_{mol} (due to the CO-to- H_2 conversion factor) and M_* (due to the stellar mass-to-light ratio) do not affect our results. We find that the inferred dynamical masses are consistent with the baryonic masses, suggesting that TDGs are nearly devoid of DM. This is illustrated in Figure 3 (left), where we compare the $M_{\text{dyn}}/M_{\text{bar}}$ ratio of TDGs with that of dwarf irregulars [15] (Irrs; blue dots from) and starburst dwarfs [16] (BCDs; black squares from). TDGs have smaller values of

$M_{\text{dyn}}/M_{\text{bar}}$ than typical dwarfs, pointing to an apparent lack of DM. In particular, we do not confirm the high values of $M_{\text{dyn}}/M_{\text{bar}}$ from [8] for the TDGs around NGC 5291. This happens because we estimate (i) lower values of V_{rot} due the technical reasons discussed above, and (ii) higher values of M_{gas} since we integrate the HI flux out to larger radii than [8] (as suggested by our 3D disc models).




Figure 2. Top panels: optical image (*left*), total HI map (*middle*), and HI velocity field (*right*). The dashed ellipse corresponds to the disc model for the assumed inclination. The cross and dashed line illustrate the kinematical centre and major axis, respectively. In the bottom-left corner, we show the linear scale (optical image) and the HI beam (total HI map and velocity field). In the total HI map, contours are at $\sim 1.5, 3.0, 4.5,$ and $6 M_{\odot} \text{ pc}^{-2}$. **Bottom panels:** PV diagrams obtained from the observed cube (*left*), model cube (*middle*), and residual cube (*right*) along the major and minor axes. Solid contours range from 2σ to 8σ in steps of 1σ . Dashed contours range from -2σ to -4σ in steps of -1σ . The horizontal and vertical lines correspond to the systemic velocity and dynamical centre, respectively.




Figure 3. **Left panel:** Mass budget within the optical radius for different types of dwarf galaxies (see [13] for details). **Right panel:** Location of TDGs on the BTFR. V_{circ} is the circular velocity after correction for pressure support. For typical galaxies, V_{circ} is measured along the flat part of the rotation curve [22,23]. For TDGs, the shape of the rotation curve is uncertain, thus V_{circ} may not correspond to the velocity along its flat part.

3. Implications for the Dark Matter Distribution

If TDGs are in dynamical equilibrium, their low values of $M_{\text{dyn}}/M_{\text{bar}}$ provide strong support to numerical simulations where tidal forces segregate halo material from disc material. They also provide an indirect but empirical confirmation to the hypothesis that DM is distributed in pressure-supported haloes. The internal dynamics of TDGs also constrains putative “dark discs” (either baryonic or non-baryonic) in the progenitor spiral galaxies, given that any kind of rotation-supported material may form tidal tails and be accreted in TDGs. For example, it has been argued that spiral galaxies should possess a rotating disc of non-baryonic DM, which is formed by the near-plane accretion of DM-dominated satellite galaxies [18]. Our results imply that either these dark discs are significantly less massive than the baryonic one or they are too dynamically hot to be accreted by TDGs.

Pfenniger *et al.* [9] proposed very cold molecular gas as an alternative to non-baryonic DM in disc galaxies. This putative dark gas would consist of small “clumpuscles” that do not emit any detectable line and are not associated with star formation. The existence of these cold H_2 clumpuscles has been advocated to explain the origin of spiral patterns and long-lived warps in the outer regions of HI discs [19,20]. Our current results do not support the existence of large numbers of cold H_2 clumpuscles. Similarly, other forms of baryonic DM in the progenitor disc galaxies (like brown dwarfs, low-mass stars, or stellar remnants) seem very unlikely. Our errors on $M_{\text{dyn}}/M_{\text{bar}}$, however, are consistent with the expectations for “standard” dark gas, such as optically thick HI or photodissociated CO in the outer regions of molecular clouds. For the Milky Way, the Planck collaboration has estimated that the mass

of dark gas may be comparable to that of CO-traced molecular gas [21]. In principle, the values of $M_{\text{dyn}}/M_{\text{bar}}$ in TDGs may be used to identify the existence of dark gas in the progenitor galaxies, but the current uncertainties on M_{dyn} do not allow such a test.

4. The Baryonic Tully-Fisher Relation and MOND

TDGs seem to systematically deviate from the BTFR (Figure 3 right). They show small rotation velocities for their baryonic masses, in line with the observed lack of a DM halo. We stress, however, that the data-points from [22,23] consider the asymptotic velocity along the flat part of the rotation curve (V_{flat}), which minimizes the scatter on the BTFR. For the TDGs in our sample, the shape of the rotation curve is uncertain, thus it remains unclear whether we are tracing V_{flat} .

The universality of the BTFR is a key prediction of MOND [24]. According to MOND, disc galaxies can deviate from the BTFR only if (i) they are out of equilibrium [25] or (ii) they are affected by the external-field effect [26] (EFE). Both circumstances may occur here. In Lelli *et al.* [13], we perform a first-order analysis of the EFE, considering the gravitational field of the progenitor galaxies. We find that the EFE can substantially decrease the velocities expected in MOND with respect to the isolated case (corresponding to the BTFR), but a systematic discrepancy persists between MOND and observations (at the level of $\sim 1-2\sigma$ for individual galaxies). We stress that our EFE analysis formally applies to the one-dimensional case and neglects the direction and time-variation of the external field. Numerical simulations are needed to investigate the EFE in detail and address the process of TDG formation and relaxation in the context of MOND.

Acknowledgments

The work of Federico Lelli and Stacy S. McGaugh is made possible through the support of a grant from the John Templeton Foundation.

Author Contributions

Federico Lelli wrote the manuscript with contributions from all authors. See Lelli *et al.* [13] for the full list of authors that contributed to this project.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Weilbacher, P.M.; Duc, P.A.; Fritze v. Alvensleben, U.; Martin, P.; Fricke, K.J. Tidal dwarf candidates in a sample of interacting galaxies. *Astron. Astrophys.* **2000**, *358*, 819–834.
2. Boquien, M.; Duc, P.A.; Galliano, F.; Braine, J.; Lisenfeld, U.; Charmandaris, V.; Appleton, P.N. Star Formation in Collision Debris: Insights from the Modeling of Their Spectral Energy Distribution. *Astrophys. J.* **2010**, *140*, 2124–2144.

3. Zwicky, F. Multiple Galaxies. In *Ergebnisse der exakten Naturwissenschaften*; Springer Berlin Heidelberg: Berlin, Germany, 1956; 344–385.
4. Bournaud, F.; Duc, P.A. From tidal dwarf galaxies to satellite galaxies. *Astron. Astrophys.* **2006**, *456*, 481–492.
5. Wetzstein, M.; Naab, T.; Burkert, A. Do dwarf galaxies form in tidal tails? *Mon. Not. R. Astron. Soc.* **2007**, *375*, 805–820.
6. Ploeckinger, S.; Recchi, S.; Hensler, G.; Kroupa, P. Chemodynamical evolution of tidal dwarf galaxies - II. The long-term evolution and influence of a tidal field. *Mon. Not. R. Astron. Soc.* **2015**, *447*, 2512–2525.
7. Barnes, J.E.; Hernquist, L. Formation of dwarf galaxies in tidal tails. *Nature* **1992**, *360*, 715–717.
8. Bournaud, F.; Duc, P.A.; Brinks, E.; Boquien, M.; Amram, P.; Lisenfeld, U.; Koribalski, S.; Walter, F.; Charmandaris, V. Missing mass in collisional debris from galaxies. *Science* **2007**, *316*, 1186–1169.
9. Pfenniger, D.; Combes, F.; Martinet, L. Is dark matter in spiral galaxies cold gas? I. Observational constraints and dynamical clues about galaxy evolution. *Astron. Astrophys.* **1994**, *285*, 79–93.
10. Milgrom, M. MOND and the Mass Discrepancies in Tidal Dwarf Galaxies. *Astrophys. J.* **2007**, *667*, L45–L48.
11. Gentile, G.; Famaey, B.; Combes, F.; Kroupa, P.; Zhao, H.S.; Tiret, O. Tidal dwarf galaxies as a test of fundamental physics. *Astron. Astrophys.* **2007**, *472*, L25–L28.
12. Kroupa, P. The Dark Matter Crisis: Falsification of the Current Standard Model of Cosmology. *Publ. Astron. Soc. Aust.* **2012**, *29*, 395–433.
13. Lelli, F.; Duc, P.A.; Brinks, E.; Bournaud, F.; McGaugh, S.S.; Lisenfeld, U.; Weilbacher, P.M.; Boquien, M.; Revaz, Y.; Braine, J.; *et al.* Gas dynamics in tidal dwarf galaxies: disc formation at $z=0$. **2015**, arXiv:1509.05404.
14. Duc, P.A.; Paudel, S.; McDermid, R.M.; Cuillandre, J.-C.; Serra, P.; Bournaud, F.; Cappellari, M.; Emsellem, E. Identification of old tidal dwarfs near early-type galaxies from deep imaging and H I observations. *Mon. Not. R. Astron. Soc.* **2014**, *440*, 1458–1469.
15. Swaters, R.A.; Sancisi, R.; van Albada, T.S.; van der Hulst, J.M. The rotation curves shapes of late-type dwarf galaxies. *Astron. Astrophys.* **2009**, *493*, 871–892.
16. Lelli, F.; Verheijen, M.; Fraternali, F. Dynamics of starbursting dwarf galaxies. III. A H I study of 18 nearby objects. *Astron. Astrophys.* **2014**, *566*, A71.
17. Lynden-Bell, D. Statistical mechanics of violent relaxation in stellar systems. *Mon. Not. R. Astron. Soc.* **1967**, *136*, 101.
18. Read, J.I.; Lake, G.; Agertz, O.; Debattista, V.P. Thin, thick and dark discs in Λ CDM. *Mon. Not. R. Astron. Soc.* **2008**, *389*, 1041–1057.
19. Revaz, Y.; Pfenniger, D. Bending instabilities at the origin of persistent warps: A new constraint on dark matter halos. *Astron. Astrophys.* **2004**, *425*, 67–76.
20. Revaz, Y.; Pfenniger, D.; Combes, F.; Bournaud, F. Simulations of galactic disks including a dark baryonic component. *Astron. Astrophys.* **2009**, *501*, 171–187.
21. Planck Collaboration. Planck early results. XIX. All-sky temperature and dust optical depth from Planck and IRAS. Constraints on the "dark gas" in our Galaxy. *Astron. Astrophys.* **2011**, *536*, A19.

22. McGaugh, S.S. The Baryonic Tully-Fisher Relation of Galaxies with Extended Rotation Curves and the Stellar Mass of Rotating Galaxies. *Astrophys. J.* **2005**, *632*, 859–871.
23. McGaugh, S.S. The Baryonic Tully-Fisher Relation of Gas-rich Galaxies as a Test of Λ CDM and MOND. *Astrophys. J.* **2012**, *143*, 40.
24. Milgrom, M. A Modification of the Newtonian Dynamics—Implications for Galaxy Systems. *Astrophys. J.* **1983**, *270*, 384.
25. McGaugh, S.S.; Wolf, J. Local Group Dwarf Spheroidals: Correlated Deviations from the Baryonic Tully-Fisher Relation. *Astrophys. J.* **2010**, *722*, 248–261.
26. Bekenstein, J.; Milgrom, M. Does the missing mass problem signal the breakdown of Newtonian gravity? *Astrophys. J.* **1984**, *286*, 7–14.

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).